


BOLETÍN

DEL

MUSEO NACIONAL DE HISTORIA NATURAL DEL PARAGUAY



Boletín del Museo Nacional de Historia Natural del Paraguay

(ISSN 1680-4031)

El Boletín del Museo Nacional de Historia Natural del Paraguay se publica en un volumen y dos números por año. Publica trabajos originales sobre aspectos varios en las áreas de Botánica, Zoología, Paleontología y Geología Descriptiva, cubriendo la Región Neotropical, principalmente Paraguay y regiones limítrofes.

EDITOR: Bolívar R. Garcete-Barrett

CORREO ELECTRÓNICO: bolosphex@sce.cnc.una.py

EDITOR ASOCIADO: Emilio Buongermini

CORREO ELECTRÓNICO: ebuongermini@paraguaysilvestre.org.py

COMITÉ REVISOR

Los editores agradecen de manera especial a los expertos citados a continuación, por su asistencia en la revisión crítica de diferentes manuscritos.

Sebastián Apesteguía (Museo Argentino de Ciencias Naturales, Argentina)

Norman Scott (Natural History Museum of Los Angeles County, U.S.A.)

James M. Carpenter (American Museum of Natural History, U.S.A.)

Junichi Kojima (Ibaraki University, Japón)

John A. Kochalka (Museo Nacional de Historia Natural del Paraguay, Paraguay)

Alberto Yanosky (Fundación Guyra Paraguay, Paraguay)

Daniel V. Hagan (Georgia Southern University, U.S.A.)

Tomás Ríos (Museo Nacional de Historia Natural del Paraguay, Paraguay)

Marizza Quintana (Museo Nacional de Historia Natural del Paraguay, Paraguay)

Carolina Pedroso (Centro de Datos para la Conservación, Paraguay)

Rocío Botta (Centro de Datos para la Conservación, Paraguay)

John P. T. Boorman, (retirado, Guildford, U.K.)

Las opiniones vertidas en los artículos son entera responsabilidad de los respectivos autores.

Museo Nacional de Historia Natural del Paraguay



Dirección de Investigación Biológica
Dirección General de Protección y Conservación de la Biodiversidad
Secretaría del Ambiente
Presidencia de la República



DIRECCIÓN: Sucursal 1 Campus U.N.A., 2169 CDP, Central XI, San Lorenzo, PARAGUAY

TELEFAX: 011-595-21-585208

CORREO ELECTRÓNICO: mnhnp@sce.cnc.una.py

SITIO EN INTERNET: www.mnhnp.org

Esta publicación fue financiada por el proyecto PAR98/G33



Primer semestre del año 2003. Se imprimieron 500 ejemplares

Ilustración de la portada: Perfiles columnares de algunas localidades con los sedimentos de relleno del rift de Asunción.

**REARING *CULICOIDES BAMBUSICOLA* (LUTZ, 1913) (DIPTERA,
CERATOPOGONIDAE) IN LABORATORY.
OBSERVATIONS AND NEW RECORDS**

MARIA M. RONDEROS¹ and FLORENTINA DIAZ²

^{1,2}Departamento Científico de Entomología, Facultad de Ciencias Naturales y Museo.
Paseo del Bosque s/n, 1900 La Plata, Argentina. ¹e-mail: ronderos@museo.fcnym.unlp.edu.ar

Abstract.- *Culicoides bambusicola* Lutz, 1913 biting midges (Diptera, Ceratopogonidae) has immature stages that breed in tacuara cane internodes, and in artificial microholders. They were reared from larvae I to adult in petri dishes on substrate containing water from their natural environment. They exhibit the ultrastructural characters of *Culicoides* larvae that breed in tree holes. In this paper reports on larval growth, feeding behavior, locomotion behavior and the duration of immature stages are provided. The technique has significant advantages for small colony maintenance in the laboratory and it allows permanent observation of developing immatures in experimental studies. Besides, this species is recorded for the first time in the Province of Buenos Aires (Argentina).

Resumen.- *Culicoides bambusicola* Lutz, 1913 (Diptera, Ceratopogonidae) presenta los estados inmaduros que crían en los internodos de cañas de tacuara y microcontenedores artificiales. Fueron criados desde larva I a adultos en pequeñas cápsulas de petri utilizando como sustrato el agua de su ambiente natural. La larva de *Culicoides bambusicola* exhibe los caracteres ultraestructurales de las larvas de *Culicoides* que crían en huecos de árboles. En este trabajo se da a conocer el desarrollo larval, comportamiento alimenticio, locomoción y duración de los estados inmaduros. La técnica demuestra ventajas significativas sobre el mantenimiento de colonias pequeñas en laboratorio y permitió la permanente observación del desarrollo de los estados preimaginales en estudios experimentales. Además, se da a conocer por primera vez el registro de esta especie para la Provincia de Buenos Aires (Argentina).

Scientific knowledge concerning adults of biting midges, *Culicoides* Latreille, is advanced in comparison to what is known of immature stages. Despite the interest in the genus, there is comparatively little knowledge about them. In most cases, the larval habitats are largely unknown, and details concerning to larval biology and feeding behavior remained obscure. The accumulation of data and observations during the past years have contributed significantly to the knowledge of the genus. Most information on breeding sites is based on studies on North American species: Thomsen's (1937), Jamback (1965), Battle & Turner (1971, 1972), Jones (1961), Blanton & Wirth (1979), Knausenberger (1986 dissert.), Mullen & Hribar (1988), and more recently, Hribar & Mullen (1991).

The larval biology of Neotropical *Culicoides* biting midges while receiving considerable study, remains for the most part unknown. Wirth & Blanton (1959) described the breeding sites for only 15 out of the 88 reported

Panamanian species of *Culicoides*, some of which are biologically closely associated to the local flora, in which the immature stages are found in plants, plant products, or in the remaining water associated with them. Williams (1964) made a significant contribution to the biology of Neotropical *Culicoides*, reporting the larval habitats for 24 species in Trinidad, from which 15 species were found associated with plant materials. Clastrier (1971) reared one species from moss growing on the palm trunk and from moist material accumulated in the axils of palm leaves in French Guinea. Vitale *et al.* (1981) reported results of a study on rearing from arboreal habitats in Panama.

There have been relatively few morphological studies of larval *Culicoides*, especially for the neotropical fauna (being described only the larvae of 12 of the 271 known species, Borkent & Spinelli (2000). The major work is on North American species. It was done by Murphee & Mullen (1991), but little is known about the ultrastructure of the larvae of biting midges, in

relation to their biology. Later, Hribar (1993) discussed the relationship between feeding behavior and morphology of mouthparts.

The larva and pupa of *Culicoides bambusicola* have recently been redescribed by Ronderos & Spinelli (2000), from material collected in the area influenced by the Yacyreta dam lake in the Paraná River (Argentina-Paraguay).

The purpose of this paper is to provide data on the biology traits: breeding location, locomotion and feeding behavior of the larvae *Culicoides bambusicola*., we describe the technique for rearing in the lab, and we also present here the first record of this species in the Province of Buenos Aires.

MATERIALS AND METHODS

A total (colony) of 108 larvae, and 8 adults of *Culicoides bambusicola* were used in this study. They were collected in sites where sampled, from tacuara internodes (Argentina, Corrientes, Garapé, 17 VI 1999; Corpus, Pto. Maní, V 2000; II 2001), and in artificial microholders (Argentina, Buenos Aires). Standing water and/or moist organic matter were removed from tacuara and were aspirated through pipettes and carried to the laboratory in tubes containing water from their natural environment. In laboratory, they were placed in styrofoam cups and covered with glass lids to prevent adults from escaping. They were observed daily to record different phases of development, and water containing nutrients, Oligochaeta (immature stages of Tubicidae) and detritus were added, and observed daily for pupation. The pupae were extracted from samples by micropipette, and placed individually in small cotton-stoppered test tubes with ca. 5 ml of water. The tubes were checked daily for emergence of adults.

The remaining sample was transferred. Larvae were cleared in a saturated solution of phenol in ethanol. The some ethanol-preserved specimens were cleared in 10 percent K(OH) for 24-48 hours. These larvae were mounted

on microscope slides in Canada balsam Wirth & Marston (1968). Larvae and pupae were also examined using scanning electron microscopy (SEM). For that purpose, they were mounted by using the techniques of Ronderos *et al.* (2000).

All emerging arthropods were collected and stored individually in vials of 70% alcohol for later observation, by means of an isolating individual rearing method association of larval skin and pupal case with a specific adult specimen. It also permitted emergence of a large number of adults from the less-disturbed, nutrient-containing original sample.

The material examined was deposited in the collection of the Museo de La Plata, Argentina in 70% alcohol in hemolysis tubes, some others on microscope slides, and on SEM stubs.

The description, illustration and SEM microphotographs, of the larva, pupa and adult of *Culicoides bambusicola* can be seen in Ronderos & Spinelli (2000).

RESULTS

Larval habitats: *Culicoides bambusicola* larvae were collected from holes of tacuara cane *Guada trini* (Ness) Ness et Rupr, at different heights, and in closed and open tacuara, and in very small artificial microholders in tree holes. This is the first species from the *cotocripus* species group found where breeding. Generally *Culicoides* larvae occur in many aquatic and semiaquatic habitats Mullen & Hribar (1988), *Culicoides* genus is also adapted to utilizing smaller ecological habitats. Therefore it is difficult to identify breeding sites, and to collect and associate larvae with adults Hynes (1984). The new record in Gonnet was collected with *Dasyhelea necrophila* Spinelli & Rodriguez, 1999 (Ronderos *et al.* In press).

In some cases, associations are evident between certain taxonomic structures and larval habitats. Ronderos & Spinelli (2000) mention that *Culicoides bambusicola* exhibits: the same stated characters by Murphree & Mullen (1991) for the larvae of tree hole breeding spe-

cies: diffuse yellowish brown thoracic pigmentation, the extensive epipharyngeal lateral curtains, and large setae of the caudal segment. The greatest variation is in OLR (ratio of caudal segment-length to length of setae "o") and average of caudal setae. In particular, the average on caudal setae "ö" and "ĩ" differs considerably from typical average.

Larval food resources: *C. bambusicola* feed from Oligochaeta (immature stages of the Tubicidae family). These larvae have long, narrow heads with the mouthparts directed anteriorly, a sclerotized labium, present a lightly built pharynx and hooked mandible, broad and pointed, it has smooth and finely serrate hypostomata single-toothed mandibles. Significant variation in form and arrangement of the palatal sensilla and appendages exists among *Culicoides bambusicola* in relation to feeding. *C. bambusicola* was fed by a supply, of small aquatic organisms as potential prey. Tissues were removed from the mandibles and taken out from its exoskeleton. Movements of the epipharynx cause the suction inside the cibarium and from the stomodeum by peristalsis.

Larval locomotion: *C. bambusicola* shows a good adaptation for swimming and typical active movement, the young larvae (I-II) moves much more slowly, but the later instar (III-IV) exhibits serpentine locomotion. Larvae are positively phototrophic, their activity notably grows when lay under the ME light and their positive phototropismo come down into term to pupae state (notable difference in relation to larvae *Culicoides insignis* Lutz, 1913) (Ronderos personal observation, 2000).

Developmental times: Adult phase is very short (5-10 days) comparing to larval phase lifetime that varies between 30-35 days. The first larval instar is the longest in life cycle of *C. bambusicola*, Its development requires two or three weeks, but from the fourth larval instar to pupation requires one week. *C. bambusicola* completes its development in as little as six days when placed in cotton-stop-

pered vials individually, from two to three days when reared at 27 ° C in the same place. Larval cycle lasted 30 days more or less in the lab.

Therefore, larval lifetime represents more than 80% in biotic cycle of this species without egg time.

New Records: *Culicoides bambusicola* (Lutz): Argentina, Buenos Aires, Gonnet, XII-2000, Ronderos-Baran, 1 larva; City Bell, 6-III-2001, 2 larvae, col. Ronderos, reared in laboratory.

DISCUSSION

This work tries to find a solution to the relative difficulty in locating developmental sites or collecting and associating adults. Factors which have contributed to difficult collecting are: small size (3-6 mm), colour (whitish), difficulty in identifying breeding sites and in collecting and associating larvae to adults Hynes (1984).

However, significant progress has been made in recent years in the larval taxonomy and habitat associations. The few *Culicoides* larvae described in detail from tree holes *C. hoffmani* Fox, 1946, *C. paraensis* (Goeldi), 1905 and *C. furens* (Poey, 1851) are unique in possessing four pairs of long perianal bristles Linley & Kettle (1960) and Murphree & Mullen (1991). These are present in many other Ceratopogonid genera but have not been recorded in many species of *Culicoides*. It would be interesting to find out whether other tree-hole breeding species have similar perianal bristles.

From a morphobehavioral perspective this variation is most evident in the mandibles, epipharynx and hypostoma. Larvae that poses a heavily sclerotized epipharynx also poses a heavily sclerotized hypopharynx.

One of the least known aspects of *Culicoides* larvae is the nature of their food and feeding behavior. Direct feeding observations have been made for only a few representative species. The species whose larvae have lightly built pharynges may feed on algae

(*Culicoides* sp.), or on detritus, or conversely they are carnivorous (*C. furens* can be reared on free-living nematodes) (Bildlingmayer, personal observation). Variety in larval feeding shows variety in feeding habitat. It were observed oligochaetes, fungi and non-identifiable particles in the alimentary tract with optical microscope oil immersion (100x) with phase contrast. Other authors (Mullens & Velten, 1994; Breidenbaugh & Mullens, 1999) offered several food sources, including a rich nutrient liquid diet consisting of bacteria, algae and added nematodes.

This work indicates that *C. bambusicola* feeds exclusively on microorganisms carried in the standing water and organic matter of their natural environment.

Regarding the locomotion, *Culicoides* larvae are generally considered to be good swimmers, although detailed observations have been made for only a few species. Linley (1986) has demonstrated that swimming ability is highly influenced by the instar and size of larvae, water temperature, and viscosity of the medium. *C. bambusicola* larvae use their specialized posterior setae for their strong activity when swimming; the reason may reside in the cleanliness and tranquility of water in the tree holes.

Development times for *Culicoides* larvae are highly variable depending on the species, temperature, water quality, and availability of qualified food.

This bloodsucking genus *Culicoides* is an important pest for man and domestic animals, but the information on the life cycle and taxonomy of the immature stages of *Culicoides* is so limited that progress on its knowledge is very limited.

Furthermore, it would be important to remark that two light traps set during two years near tacuara canes in the sampling stations, did not collect specimens of *C. bambusicola*, suggesting that the species is poorly attracted by light. The previous valid record for *C. bambusicola* is also based on specimens collected in cane internodes.

This is one of the first studies of the breeding sites of the different species of the genus, and the relationship existing with the taxonomic ultrastructure. Further taxonomic work with the immature stages will, no doubt enhance the prospects for more detailed characterizations of the larval biology and breeding, locomotion and feeding behavior of *Culicoides* species in the future.

This record of *C. bambusicola* in Gonnet was collected with other Ceratopogonidae: *Dasyhelea necrophila* Spinelli & Rodriguez (1999). The larva of this species exhibits the character states typical of "herbivorous" ceratopogonids (Ronderos *et al.*, in press). Precisely specific adaptations to different larval habitats will allow researchers to set limits on the actual distribution of certain species, since immature states can show different morphoecological types.

ACKNOWLEDGMENTS

We gratefully acknowledge the critiques and helpful suggestions by Dra. María Marta Cigliano, Dr. Gustavo Spinelli.

REFERENCES

- Battle, F. V. & Turner, e. C. 1971. The insects of Virginia. N° 3. A systematic review of genus *Culicoides* (Diptera: Ceratopogonidae) of Virginia, with a geographic catalog of the species occurring in the eastern United States north of Florida. Virginia Polytech. Inst. State. Univ. Res. Div. Bull., 44: 1-129.
- Battle, F. V. & Turner, E. C. 1972. Some nutritional and chemical properties of the larval habitats of certain species of *Culicoides* (Diptera: Ceratopogonidae). *J. Med. Entomol.*, 9: 32-35.
- Blanton, F. S. & Wirth., W. W. 1979. The sand flies (*Culicoides*) of Florida (Diptera: Ceratopogonidae). *Arthropods of Florida and Neighboring Land Areas*, 10: 1-204.
- Borkent, A. & Spinelli, G. R. 2000. Catalog

- of the New World biting midges south of the United States of America (Diptera: Ceratopogonidae). *Contributions on Entomology, International*, 4 (1): 1-107.
- Breindenbaugh, M. S. & Mullens, B. A. 1999. Two new western Nearctic *Culicoides* Latreille (Diptera: Ceratopogonidae) described from all stages. *Proc. Ent. Wash*, 10 (1): 149-163.
- Clastrier, J. 1971. Deux nouveaux *Culicoides* (Diptera: Ceratopogonidae) Guyane Francaise. *Ann. Parasit. Hum. Comp.*, 46: 285-294.
- Hribar, L. J. 1993. Mouthpart Morphology and Feeding Behavior of Biting Midge Larvae (Diptera: Ceratopogonidae). In: *Functional Morphology of Insect Feeding*. C. W. SCHAEFER & R. A. B. LESCHEN, Eds., págs. 43-58. Thomas Say Publications in Entomology, Proceedings of the Entomological Society of America. Lanham, MD.
- Hribar, L. J. & Mullen, G. R. 1991. Comparative morphology of the mouthparts and associated feeding structure of biting larvae (Diptera: Ceratopogonidae). *Contrib. Amer. Entomol. Inst.*, 26: 1-71.
- Hynes, H. B. N. 1984. The relationship between the taxonomy and ecology of aquatic insects. In: *The ecology of aquatic insects*. V. H. RESH & D. M. ROSEMBERG, Eds., págs. 9-23. Praeger Sci. Publ., New York.
- Jamback, H. 1965. The *Culicoides* of the New York State (Diptera: Ceratopogonidae). *Bull. N.Y. St. Mus. Sci. Serv.*, 399: 1-154.
- Jones, R. H. 1961. Observation on the larval habitats of some North American species of *Culicoides* (Diptera: Ceratopogonidae). *Ann. Entomol. Soc. Am.*, 54: 702-710.
- Knausenberger, W. L. 1986. Contribution to the autoecology and ecosystematics of immature Ceratopogonidae (Diptera), with emphasis in the tribes Heteromyiini and Sphaeromyiini in the Middle Atlantic United States. Ph.D. Dissert. Virginia Polytech. State Univ. Blacksburg. VA, 728 págs.
- Linley, J. R. 1986. Swimming behavior of the larva of *Culicoides variipennis* (Diptera: Ceratopogonidae) and its relationships to temperature and viscosity. *J. Med. Entomol.*, 23: 473-483.
- Linley, J. R. & Kettle, D. S. A. 1960. Description of the larva of *Culicoides hoffmani* Fox (Diptera: Ceratopogonidae) and some observations on its feeding habits. Rept. to Ministry of Health, Jamaica, W. I.
- Mullen, G. R. & Hribar, L. J. 1988. Biology and feeding behavior of Ceratopogonidae larvae (Diptera: Ceratopogonidae) in North America. *Bull. Soc. Vector Ecol.*, 13: 60-81.
- Mullens, B. A. & Velten, R. K. 1994. Rearing *Culicoides variipennis sonoriensis* (Diptera: Ceratopogonidae) on agar and nematodes. *Journal of Med. Entomol.*, 31: 175-177.
- Murphree, C. S. & Mullen, G. R. 1991. Comparative larval morphology of the genus *Culicoides* Latreille (Diptera: Ceratopogonidae) in North America with a key to species. *Bull. Soc. Vec. Ecol.*, 16 (2): 269-399.
- Ronderos, M. M. & Spinelli, G. R. 2000. The larvae and pupa of *Culicoides bambusicola* Lutz observed with SEM, and additional notes on the adults (Diptera: Ceratopogonidae). *Trans. Amer. Entomol. Soc.*, 126 (1): 133-144.
- Ronderos, M. M., Spinelli, G. R. & Sarmiento, P. 2000. Preparation and mounting of biting midges of the genus *Culicoides* Latreille (Diptera: Ceratopogonidae) to be observed with a scanning electron microscope. *Transactions of the American Entomological Society*, 126 (1): 125-132.
- Ronderos, M. M., Spinelli, G. R., Huerta, H.

- & Diaz, F. (in press). Immature stages of two species of *Dasyhelea* Kieffer, 1991 (Diptera: Ceratopogonidae).
- Spinelli, G. R. & Rodriguez, E. A. 1999. A new species of *Dasyhelea* from Argentina described as adult, pupae and fourth instar larvae (Diptera: Ceratopogonidae). *Neotropica*, 45: 59-62.
- Thomsen's, L. C. 1937. Aquatic Diptera. Part V, Ceratopogonidae. Mem. Cornell Univ. *Agric. Expt. Stn.*, 210: 57-80.
- Vitale, G. C., Wirth, W. W. & Aitken, T. H. G. 1981. New species and records of *Culicoides* reared from arboreal habitats in Panamá, with a synopsis of the *debilipalpis* group (Diptera: Ceratopogonidae). *Proc. Ent. Soc. Wash.*, 83: 140-159.
- Wirth, W. W. & Blanton, F. S. 1959. Biting midges for the genus *Culicoides* from Panamá (Diptera: Heleidae). *Proc. U.S. Nat. Mus.*, 109: 237-482.
- Wirth, W. W. & Marston, N. 1968. A method for mounting small insects on microscope slides in Canada balsam. *Annals of the Entomological Society of America*, 61: 783-784.
- Williams, R. W. 1964. Observations on habitats of *Culicoides* larvae in Trinidad (Diptera: Ceratopogonidae). *Ann. Ent. Soc. Am.*, 57: 462-466.

INSTRUCCIONES A LOS AUTORES

El Boletín del Museo Nacional de Historia Natural del Paraguay se publica en un volumen por año, dividido en dos números. Las fechas límite para recepción de manuscritos son 1 de marzo para la edición de mayo (n° 1) y 1 de setiembre para la edición de noviembre (n° 2). En caso de no tener un número completo para la edición de mayo, se publicará un volumen de doble número en noviembre.

Se aceptan trabajos de investigación originales diversos en las áreas de Botánica, Zoología, Paleontología y Geología Descriptiva, cubriendo la Región Neotropical y preferentemente el Paraguay y regiones limítrofes. Se aceptan trabajos en Español o en Inglés.

Los manuscritos deben presentarse en archivo electrónico generado en Microsoft Word y copia impresa en papel tamaño carta con todos los márgenes de 2,5 cm y texto en fuente Times New Roman tamaño 11. No se aceptarán pies de página.

Las figuras deben ser originales, con número de referencia escrito a lápiz al dorso o en su defecto archivos electrónicos numerados, de buena resolución en formatos JPG, TIF o PNG. Los pies de ilustración deben ir en hoja aparte, indicando claramente los números de referencia de las ilustraciones originales o los archivos respectivos. Las tablas deben ir impresas por separado y en versión electrónica en archivo generado en Microsoft Excel.

Toda la documentación relacionada con el artículo debe enviarse al Editor del Boletín, Museo Nacional de Historia Natural del Paraguay, Sucursal 1 Campus U.N.A, 2169 CDP, Central XI, San Lorenzo, Paraguay. Los archivos electrónicos pueden enviarse a la dirección electrónica vigente del museo o del editor. Se pide que los autores provean nombre, dirección postal y correo electrónico de al menos dos revisores potenciales.

La primera página del manuscrito debe llevar los siguientes datos: **1)** título conciso e informativo en letra mayúscula, **2)** nombre del autor o autores, **3)** dirección completa del autor o autores (incluyendo dirección electrónica si existe), **4)** resumen en español y **5)** abstract en inglés.

El cuerpo del manuscrito puede constar de las siguientes partes ordenadas, cada una titulada en letra mayúscula: **1)** Introducción, **2)** Materiales y Metodología, **3)** Resultados y Discusión, **4)** Conclusión y **5)** Literatura. Se aceptan modificaciones de este esquema siempre que sigan una secuencia lógica equivalente a lo propuesto.

Los trabajos deberán respetar las disposiciones de los códigos de nomenclatura Zoológica y Botánica vigentes. Los nombres científicos deben escribirse en *itálicas*. Se sugiere que los nombres científicos se escriban completos, incluyendo autor(es), al menos la primera vez que se mencionan. La citación de autores de nombres científicos es obligatoria en trabajos de índole taxonómico. Los nombres genéricos al principio de una oración deben escribirse completos.

Las citas bibliográficas deberán hacerse de acuerdo a los siguientes ejemplos: López (1992) o (López, 1992). Cuando un trabajo tiene dos autores se mencionarán ambos apellidos y cuando sean más se citará como en los ejemplos: López *et al.* (1991) o (López *et al.*, 1991).

En la sección Literatura se deben incluir los trabajos citados en el manuscrito o que merecen mención justificada por su importancia en el tema tratado. Las referencias deben ir por orden alfabético y cronológico y cada una siguiendo el modelo de secuencia: Autor. Año. Título. Publicación serial o Casa editora, Volumen (Número) : Total de páginas o Secuencia de páginas. Abajo hay algunos ejemplos:

Carpenter, J.M. 1986. A synonymic generic checklist of the Eumeninae (Hymenoptera: Vespidae). *Psyche*, 93 (1 – 2) : 61 – 90.

Carpenter, J.M. & J. Vecht. 1991. A study of the Vespidae described by William J. Fox (Insecta: Hymenoptera), with assessments of taxonomic implications. *Annals of Carnegie Museum*, 60 (3) : 211 – 241.

Polazek, A., S. Abd-Rabou & J. Huang. 1999. The Egyptian species of *Encarsia* (Hymenoptera: Aphelinidae); a preliminary review. *Zoologische medelingen Leiden*, 73 : 131 – 163.

Hanson, P. & A.S. Menke. 1995. The sphecid wasps (Sphecidae). Capítulo 17, pp. 621 – 646, *en* Hanson P. & I.D. Gauld (editores). *The Hymenoptera of Costa Rica*. Oxford Science Publications/The Natural History Museum, London. 893 pp.

Richards, O.W. 1978. The social wasps of the Americas excluding the Vespinae. *British Museum (Natural History)*, London. 580 pp.

INSTRUCTIONS TO AUTHORS

Boletín del Museo Nacional de Historia Natural del Paraguay is published a volume a year, divided in two numbers. Deadline for manuscript reception is March 1 for the May edition (n° 1) and September 1 for the November edition (n° 2). A double number volume will be published in November if no papers were available to complete de May edition.

The editorial accepts original research papers on several aspects of Botany, Zoology, Paleontology and Descriptive Geology, covering the Neotropical Region, preferably Paraguay and neighbouring areas. Papers wrote in Spanish or English will be accepted.

The manuscripts should be submitted as electronic files in Microsoft Word format and printed in letter size paper with 25 mm margins and text in Times New Roman font, size 11. Footnotes will not be accepted.

Figures should be submitted as original hard copies, with reference numbers penciled on back or, alternatively as numbered electronic files with good resolution in JPG, TIF or PNG format. The figure legends must go in a separate page, clearly indicating the reference numbers of the original illustrations or files. Tables should be printed separately and electronic files containing them should be made in Microsoft Excel format.

All the documentation related to the manuscript must be sent to: Editor del Boletín, Museo Nacional de Historia Natural del Paraguay, Sucursal 1 Campus U.N.A, 2169 CDP, Central XI, San Lorenzo, Paraguay. The electronic files could be sent to the available e-mail address of the museum or of the editor. Authors are asked to provide name, address and e-mail of at least two potential referees.

The first page of the manuscript must contain the following data: **1)** short and concise title in capitals, **2)** name of the author(s), **3)** complete address of the author(s) (including e-mail address if available), **4)** Spanish 'resumen' and **5)** English abstract.

The manuscript body could be composed by the following ordered parts, each one entitled in capitals: **1)** Introduction, **2)** Materials and Methods, **3)** Results and Discussion, **4)** Conclusion and **5)** Literature. Modifications could be accepted if they follow a logic sequence equivalent to the one here proposed.

Papers must respect the rules of the codes on Zoology and Botany in force. Scientific names must be in *italics*. It is suggested that scientific names should be mentioned complete, including author(s) at least in the first mention. Authority is mandatory in taxonomic papers. Generic names must be completely spelled at the beginning of a sentence.

References in the text should follow the examples: López (1992), or (López, 1992). Papers with two authors should mention both names, and papers with more authors should follow the examples: López *et al.* (1991), or (López *et al.*, 1991).

The Literature section must include all the works referred in the text and could include those with justified influence on the subject. References should go in alphabetic and chronologic order, each one according to the following model: Author. Year. Title. Serial publication or editorial, Volume (Number) : Page total or page sequence. Examples are given bellow:

- Carpenter, J.M. 1986. A synonymic generic checklist of the Eumeninae (Hymenoptera: Vespidae). *Psyche*, 93 (1 – 2) : 61 – 90.
- Carpenter, J.M. & J. Vecht. 1991. A study of the Vespidae described by William J. Fox (Insecta: Hymenoptera), with assessments of taxonomic implications. *Annals of Carnegie Museum*, 60 (3) : 211 – 241.
- Polazek, A., S. Abd-Rabou & J. Huang. 1999. The Egyptian species of *Encarsia* (Hymenoptera: Aphelinidae); a preliminary review. *Zoologische medelingen Leiden*, 73 : 131 – 163.
- Hanson, P. & A.S. Menke. 1995. The sphecid wasps (Sphecidae). Capítulo 17, pp. 621 – 646, *en* Hanson P. & I.D. Gauld (editores). *The Hymenoptera of Costa Rica*. Oxford Science Publications/The Natural History Museum, London. 893 pp.
- Richards, O.W. 1978. The social wasps of the Americas excluding the Vespinae. *British Museum (Natural History)*, London. 580 pp.

CONTENIDO

- Wild, A. L.** The genus *Pachycondyla* (Hymenoptera: Formicidae) in Paraguay 1 - 18
- Carpenter, J. M.** Return to the subspecies concept in the genus *Zeta* (Hymenoptera: Vespidae; Eumeninae) 19 - 24
- Garcete-Barrett, B. R.** Notes on neotropical Eumeninae (Hymenoptera: Vespidae) II, the genus *Incodynerus* Willink 25 - 29
- Oliva, A.** Especies de Berosinae (Coleoptera: Hydrophilidae) de Paraguay 30 - 35
- Garcete-Barrett, B. R.** Notas sobre Eumeninae neotropicales III (Hymenoptera: Vespidae) 36 - 42
- Ronderos M. M. and F. Diaz.** Rearing *Culicoides bambusicola* (Lutz, 1913) (Diptera: Ceratopogonidae) in laboratory. Observations and new records 43 - 48
- Garcete-Barrett, B. R.** A new species of *Zethus* (Hymenoptera: Vespidae: Eumeninae) from Eastern Paraguay 49 - 51
- Carpenter, J. M. and B. R. Garcete-Barrett.** A key to the neotropical genera of Eumeninae (Hymenoptera: Vespidae) 52 - 73
- Morales, C. y C. Vitale.** Tendencias poblacionales del *Nandayus nenday* (Vieillot) (Aves: Psittacidae) en el Bajo Chaco paraguayo 74 - 79
- Garcete-Barrett, B. R.** La taxonomía del género *Cephalastor* Soika (Hymenoptera: Vespidae: Eumeninae), parte II 80 - 97
- Filippi Amábile, V. y J. L. Báez Presser.** Correlación de sedimentos de relleno del rift de Asunción en el Valle de Acahay 98 - 105